

A NEW APPROACH TO DEFORMING TRIANGULATED CATEGORIES

Alessandro Lehmann
SISSA and University of Antwerp

Motivation: the curvature problem

Let A be a dg-algebra over a field k . It is a well known problem in deformation theory [KL09] that (first order) deformations of A as a dg-algebra do not suffice to span its whole second Hochschild cohomology. It turns out [Low08; Leh24] that $\mathrm{HH}^2(A)$ parametrizes first order deformations of A as a *curved* dg algebra. This is a significant issue, since cdg algebras do not have classical derived categories [Pos10]; in particular, there is no obvious deformation of $D(A)$ corresponding to a curved deformation of A [KLN10].

The same problem was observed by Lurie in a different setting: he essentially showed in [Lur11] that the functor

$$\mathrm{Def}_A: \mathrm{d}\mathrm{g}\mathrm{art}_k \rightarrow \mathrm{Set}$$

which associates to a (dg) local artinian k -algebra R the set of R -deformations of A as a dg-algebra is not a (derived) deformation functor.

The main question that we aim to answer is the following:

Which deformation of $D(A)$ corresponds to a curved deformation of A ?

Lurking behind this one is a more fundamental question: *which notion* of deformation of a triangulated category allows for the question above to have a positive answer? The usual one – which roughly corresponds to reducing the hom-sets of an opportune resolution – cannot work: to a curved deformation of A corresponds a curved deformation of the dg-category $D(A)$, which does not have an underlying triangulated category. This question is a crucial step towards obtaining a satisfactory deformation theory for noncommutative schemes.

cdg algebras

A cdg algebra \mathcal{A} over a commutative ring R is a triple $(\mathcal{A}^\#, d_{\mathcal{A}}, c)$ where:

- $\mathcal{A}^\#$ is a graded R -algebra;
- $d_{\mathcal{A}}: \mathcal{A}^\# \rightarrow \mathcal{A}^\#$ is a degree 1 derivation;
- $c \in \mathcal{A}^\#$ is a degree 2 element such that $d_{\mathcal{A}}c = 0$ and $d_{\mathcal{A}}^2 = [c, -]$.

A cdg module M over a cdg algebra \mathcal{A} is a pair $(M^\#, d_M)$ where $M^\#$ is a graded $\mathcal{A}^\#$ -module and

$$d_M: M^\# \rightarrow M^\#$$

is a degree 1 derivation such that $d_M^2 m = cm$.

Key point: if M and N are cdg \mathcal{A} -modules, then

$$\mathrm{Hom}_{\mathcal{A}}(M, N)$$

is a *complex*, so $\mathcal{A}\text{-Mod}$ is a (pretriangulated) dg-category.

However, cdg modules have no cohomology, so there is no obvious notion of a derived category of \mathcal{A} [Pos10; KLN10].

The n -derived category

Let A_n be a cdg deformation of A over $k[t]/(t^{n+1})$. A cdg A_n -module M is n -acyclic if its associated graded with respect to the t -adic filtration is acyclic; this makes sense since each graded piece is a complex. The n -derived category $D^n(A_n)$ is the Verdier quotient $D^n(A_n) = H^0 A_n\text{-Mod}/n\text{-Ac}(A_n)$.

Theorem 1 ([LL24])

- The category $D^n(A_n)$ is generated by $n+1$ explicit compact objects $\Gamma_0, \dots, \Gamma_n$ and the projection $H^0 A_n\text{-Mod} \rightarrow D^n(A_n)$ admits both adjoints;
 - Calling A_i the induced deformation of order $i \leq n$, the restriction functor $A_i\text{-Mod} \rightarrow A_n\text{-Mod}$ induces a system of fully faithful embeddings
- $$D(A) = D^0(A_0) \xrightarrow{i_1} D^1(A_1) \hookrightarrow \dots \hookrightarrow D^{n-1}(A_{n-1}) \xrightarrow{i_n} D^n(A_n);$$
- The abelian category $Z^0 A_n\text{-Mod}$ admits a model structure presenting $D^n(A_n)$;

The classical case

If A_n has no curvature, one can consider the classical derived category $D(A_n)$. One sees that there are strictly more n -acyclics than acyclics; it turns out that $D^n(A_n)$ can be seen as a (partial) categorical resolution of $D(A_n)$. Indeed:

- There is an embedding
- $$D(A_n) \hookrightarrow D^n(A_n);$$
- The (dg) category $D^n(A_n)$ is smooth if and only if $D(A)$ is smooth.

$D^n(A_n)$ as a categorical extension

The embedding $D^{n-1}(A_{n-1}) \xrightarrow{i_n} D^n(A_n)$ admits both a left adjoint $\mathrm{Ker} t^n$ and a right adjoint $\mathrm{Coker} t^n$. We also have the functor $\mathrm{Im} t^n: D^n(A_n) \rightarrow D(A)$. These functors do not need to be derived, since they preserve n -acyclics.

Theorem 2 ([LL24])

There is a recollement

$$\begin{array}{ccc} & \mathrm{Ker} t^n & \\ \swarrow & \xrightarrow{\quad} & \searrow \\ D^{n-1}(A_{n-1}) & \xrightarrow{i_n} & D^n(A_n) \\ \nwarrow & \xleftarrow{\quad} & \swarrow \\ & \mathrm{Coker} t^n & \end{array} \quad \begin{array}{ccc} & \mathrm{Im} t^n & \\ \swarrow & \xrightarrow{\quad} & \searrow \\ & D(A) & \end{array}$$

In particular, the functor

$$\mathrm{Im} t^n: D^n(A_n) \rightarrow D(A)$$

induces an equivalence between the quotient $D^n(A_n)/D^{n-1}(A_{n-1})$ and $D(A)$.

Inductively, we see that $D^n(A_n)$ is obtained by gluing $n+1$ copies of $D(A)$.

Deformations of triangulated categories (WIP)

All triangulated categories and functors between them are appropriately enhanced. Let \mathcal{T} be a triangulated category. A first order predeformation of \mathcal{T} is the datum of a recollement

$$\begin{array}{ccc} & K & \\ \swarrow & \xrightarrow{\quad} & \searrow \\ \mathcal{T} & \xrightarrow{i} & \mathcal{T}_\varepsilon \\ \nwarrow & \xleftarrow{\quad} & \swarrow \\ & Q & \end{array}$$

together with two natural transformations $I \xrightarrow{\eta_1} K$ and $Q \xrightarrow{\eta_2} I$. The recollement data also induces a natural transformation $K \xrightarrow{\alpha} Q$; a predeformation is said to be a deformation if α induces a natural isomorphism $\bar{\alpha}$

$$\begin{array}{ccc} I & \xrightarrow{\eta_1} & K & \longrightarrow & C \\ & \searrow & \downarrow \alpha & \swarrow \bar{\alpha} & \\ D & \longrightarrow & Q & \xrightarrow{\eta_2} & I \end{array}$$

between the cone C of η_1 and the cocone D of η_2 . Denote with $\mathrm{Def}_{k[\varepsilon]}^{\mathrm{cat}}(\mathcal{T})$ the set of first order deformation of \mathcal{T} up to equivalence.

Main result: a commutative diagram of deformations

Theorem 3 (WIP)

Let \mathcal{T} be a triangulated category. There is a bijection

$$\mathrm{Def}_{k[\varepsilon]}^{\mathrm{cat}}(\mathcal{T}) \xleftrightarrow{\kappa} \mathrm{HH}^2(\mathcal{T})$$

between first order deformations of \mathcal{T} as a triangulated category and its second Hochschild cohomology.

The point of the bijection is that the gluing functor is the cone of the Hochschild cocycle.

Theorem 4 (WIP)

There is a commutative diagram of bijections

$$\begin{array}{ccc} \mathrm{cDef}_{k[\varepsilon]}^{\mathrm{Mor}}(A) & \longrightarrow & \mathrm{HH}^2(A) \\ D^1(-) \downarrow & & \downarrow \chi_A \\ \mathrm{Def}_{k[\varepsilon]}^{\mathrm{cat}}(D(A)) & \xrightarrow{\kappa} & \mathrm{HH}^2(D(A)) \end{array}$$

Where $\mathrm{cDef}_{k[\varepsilon]}^{\mathrm{Mor}}(A)$ is the set of curved deformations of A up to equivalence of 1-derived categories; the upper arrow was introduced in [Leh24], while χ_A is the characteristic morphism introduced in [Low08].

References

- [KL09] Bernhard Keller and Wendy Lowen. “On Hochschild Cohomology and Morita Deformations”. In: *Int. Math. Res. Not.* 2009 (2009).
- [KLN10] Bernhard Keller, Wendy Lowen, and Pedro Nicolás. “On the (non)vanishing of some “derived” categories of curved dg algebras”. In: *J. Pure Appl. Algebra* 214.7 (2010).
- [Leh24] Alessandro Lehmann. *Hochschild cohomology parametrizes curved Morita deformations*. 2024. arXiv: 2406.04945.
- [LL24] Alessandro Lehmann and Wendy Lowen. *Filtered derived categories of curved deformations*. 2024. arXiv: 2402.08660.
- [Low08] Wendy Lowen. “Hochschild cohomology, the characteristic morphism and derived deformations”. In: *Compos. Math.* 144.6 (2008).
- [Lur11] Jacob Lurie. *Derived Algebraic Geometry X: Formal Moduli Problems*. 2011.
- [Pos10] Leonid Positselski. *Two kinds of derived categories, Koszul duality, and comodule-contramodule correspondence*. Mem. Am. Math. Soc. 2010.